REFERENCES

1. Abesh Kumar, Bhardwaj Manish (2011). Diversity and indicator species of moth (Lepidotera: Heterocera) assemblages in different vegetation zones in gangotri landscape, Wetern Himalaya, India. ENVIS Bulletin: Arthropods and their Conservation in India. Vol. No. 14 (1). 114-129.
2. A.T.M.F. Islam, M.H. Islam, A.S.M. Saifullah and Akira Yamanaka (2013). A preliminary report of moth's fauna in the campus of Atomic Energy Research Establishment (AERE), Savar, Dhaka, Bangladesh. International Journal of Fauna and Biological Studies. 1 (1): 56-62.
3. Balint Horvath (2013). Diversity comparison of nocturnal macrolepidoptera communities (Lepidoptera: Macroheterocera) in different forest stands. Kaposvar. 23: 229-238.
4. D. Adiroubane and P. Kuppammal (2010). Lepidopteran fauna of Agri-Horticultural ecosystem in Karaikal region. Journal of Biopesticides. 3 (1): 1-10.
5. Dejan V. Stojanovic and Srecko B. Curcic (2011). The Diversity of Noctuid Moths (Lepidoptera: Noctuidae) in Serbia. ActaZoologicaBulgarica. Vol. 63 No. 1. 47-60.
6. D. Silhacek, C. Murphy, R.T. Arbogast (2003). Behavior and movements of Indian meal moths (*Plodiainterpunctella*hubner) during commodity infestation. Journal of Stored Products Research. Vol. No. 39. 171-184.
7. E.G. White (1991). The changing abundance of moths in Tussock grassland, 1962-1989, and 50 to 70- year trends. New Zealand Journal of Ecology. 15 (1). 5-22..
8. Gaurav Sharma (2011). Studies on lepidopterous insects associated with vegetables in Aravali Range, Rajasthan, India. Biological Forum An International Journal, 3 (1): 21-26.
9. George Mathew and V. K. Rahamathulla (1995). Biodiversity in the Western Ghats – A study with reference to moths (Lepidoptera: Heterocera) in the silent valley national park, India. Entomon, 2: 25-23.
10. Gunnar Brehm and Konrad Fiedler (2005). Diversity and community structure of geometrid moths of disturbed habitat in a montane area in the Ecuadorian Andes. Journal of Research on the Lepidoptera. 38: 1-14
11. Hari Sutrisno (2008) Moth diversity at gununghalimun- salak National Park, West Java. Journal of Biosiciences. 15(3): 111-117.
12. H.S. Rose (2001). An inventory of the moth fauna (Lepidoptera) of Jatinga, Assam, India. Zoos Print Journal. 2: 707-721.

13. Jay M. Iwasaki (2010). Moth diversity and abundance between habitats on a shade coffee plantation in Western Jamaica. Beta BetaBeta Biological Society. 81 (4): 108-113.
14. J. D. Holloway, G. Kibby and D. Peggie (2007). The families of malesian moths and butterflies. Tropical Lepidoptera. 16 (1-2): 22.
15. J. Kashefi, G.P. Markin and J. L. Littlefield (2008). Field studies of the biology of the moth Bradyrrhoagilveolella (Treitschke) (Lepidoptera: Pyralidae) as a potential biocontrol agent for chondrillajuncea. XII International Symposium on Biological Control of Weeds. 558-572.
16. J. Ramkumar, M. Swamippan, S. Raguraman and A. Sadasakthi (2010). Species diversity and seasonal abundance of fruit piercing moth complex in Tamil Nadu. Journal of Biopesticies. 3 (1): 11-15.
17. Kailash Chandra and S. Sambath (2013). Moth diversity of Tawang District, Arunachal Pradesh, India. Journal of Threatened Taxa. 5 (1): 3565-3570.
18. Kallare P. Arunkumar, Anup Kumar Sahu, AtishRanjanMohanty, Arvind K. Awasthi, Appukuttannair R. Pradeep, S. RajeUrs and JavaregowadaNagaraju (2012). Genetic diversity and population structure of Indian Golden Silkmoth (Antheraeaassama). PLOS ONE. 7 (8): 1-7.
19. KarimSaeidi (2011). Preliminary studies on natural enemies of the gypsy moth, Lymantriadispar (L.) (Lepidoptera: Lymantriidae) in Yasooj, Iran. International Journal of Agricultural Science and Soil Science. 1 (5): 151-156.
20. K. Sivasankaran, S. Gnanasekaran, D. Parandhaman and S. Ignacimutu (2011). Diverisity of Noctuid moths (Lepidoptera: noctuidae) in Tamil Nadu part of Western Ghats (Nilgiris biosphere and Kodaikanal hills), India. Elixir International Journal. 38: 4131-4134.
21. L.M. Cook (2000). Changing views on melanic moths. Biological Journal of Linnean Society. 69: 431- 441.
22. L.N. Kakati and B.C. Chutia (2009). Diversity and ecology of wild sericigenous insects in Nagaland, India. Tropical Ecology. 50 (1). 137-146.
23. Louise A. Ashton, Roger L. Kitching, Sarah C. Maunsell and Darren Bito (2011). Macrolepidopteran assemblages along an altitudinal gradient in subtropical rainforest- exploring indicators of climate change. Nature. 55 (2): 375-389.
24. Muhammad Aslam (2009). Diversity, species richness and evennesss of moth fauna of Peshawar. Pakistan Entomology. 31 (2): 99-102.

25. Muhammad Aslam (2013). Cheklist of moth fauna of Peshawar, Pakistan. Arthropods. 2 (4): 237-241.
26. Niels P. Kristensen, Malcolm J. Scoble and Ole Karsholt (2007). Lepidoptera phylogeny and systematic: the state of inventorying moth and butterfly diversity. Zootaxa. 1668: 699-747.
27. Peter Huemer (2009). Biodiversity of butterflies and moths in the national park hohetauern. 4th Symposium of the Hohe Tauern National Park for Research in Protected Areas. Confernece Vol. 135-136
28. P.S. Jagdish, P. Nirmala, M.A. Rashmi, Jayalaxmi Narayan Hedge and NeeluNangia (2009). Biology of rice moth, *Corcyra cephalonica*stainton on foxtail millet. Karnataka Journal of agricultural science. 22 (3): 674-675.
29. RaunoVaisanen (1992). Distribution and abundance of diurnal Lepidoptera on a raised bog in southern finland. Ann. Zool. Fennici. 29: 75-92.
30. ReimaLeinonen and JuhaniItamies (2000). The Finnish Environment. 485. 187-192.
31. Richard S. Peigler (2012): Diverse evidence that *Antheraeapernyi* (Lepidoptera: Saturniidae) is entirely of sericulturel origin. Trop. Lepid. Res. 22 (2): 93-99.
32. Robert Woods (2012). Brownfield sites and moth diversity in the Tees Estuary. Entomologist's Rec. J. Var. 124: 89-100.
33. Sachin A. Gurule and Santosh M. Nikam (2013). The Moth (Lepidoptera: Heterocera) of Northern Maharashtra: A Preliminary Checklist. Journal of Threatened Taxa. 5 (12): 4693-4713.
34. SamadKhaghaninia and Reza FarshbafPourabad (2009). Investigation on Biology of olive leaf worm PalpitaUnionalishb. (Lepidoptera: Pyralidae) in constant laboratory conditions. Mun. Ent. Zool. 4 (2): 320-326.
35. Tufail Ahmad and MohdShafiq Ansari (2010). Studies on seasonal abundance of diamondback moth plutellaxylostella (Lepidoptera: Yponomeutidae) on cauliflower crop. Journal of plant protection research. Vol. No. 50 (3). 280-287.
36. Viviane Glaniuppl Ferro and Helena PiccollRomanowski (2012). Diversity and composition of tiger moths (Lepidoptera: Arctiidae) in an area of Atlantic Forest in southern Brazil: is the fauna more diverse in the grassland or in the forest. Zoologia. 29 (1): 7-18
37. VedatBeskardes (2012). Lepidoptera fauna of Yuvacik dam watershed in Kocaeli, Turkey. African Journal of Agricultural Research. 7 (11): 1749-1754.

38. V. Subhalaxmi, Roger C. Kendrick, Alka Vaidya, NeelimaKalagi and AlakaBgagwat (2011). Inventory of moth fauna (Lepidoptera: Hetericera) of the Northern Western Ghats, Maharashtra, India. Journal of the Bombay Natural Histroy Society. 103 (3): 193-205.
39. Yasir Ifran Yattoo (2013). An inventory step towards species diversity of moths in pattan, District Baramulla, Jammu and Kashmir (India). AJBB Online. 3 (3): 1-4.
40. Zuhal Okyar, SelcukYurtsever, NihatAktac and GokhanCakan (2009). Some aspects of the moth (Lepidoptera, Heterocera) species diversity in Western Black Sea Region of Turkey. North-Western Journal of Zoology. 5(1): 104-120.
41. https://www.google.co.in/search?q=baramati+map&biw=1280&bih=923&tbm=isch&tbo=u&source=univ&sa=X&sqi=2&ved=0ahUKEwj9nf3Zw_nLAhVHr4MKHUflAOcQsAQIHA#tbm=isch&q=baramati+in+maharashtra+map&imgrc=2eWe8y6NZs4OVM%3A